

- Announcements
 - Recitation – Do written HW #1
 - Clickers – Count beginning today
 - Written HW due Thursday midnight
 - Make your exam reference while doing homeworks
- Homework challenges and exams
- Electrostatics
 - Types of charge
 - Tribocharging
 - Insulators and Conductors
 - Coulomb's Law
 - Superposition

iClicker

- We will start using iClicker cloud today.
- Go to www.iclicker.com and “Create an account”

Files Written_H The Expert The Expert New Tab Inbox (5,29) Launch Me Inbox (5,29) WSO2 Ider Assignmen Inbox Index of /- Advisee Li iClicker: St New Mexic Brave Help iClicke x + -

Getting Started https://ep3guide... LxArchive - by Ea... Index of /media/... ISS: ASIM - eoPor... Have You Seen B... Building a Raspb... LOFAR Shared Data Phys334 Synchronizing nt... Home Page Wav... Home Dart Leader Velo...

iClicker a mind on learning company Solutions Getting Started with iClicker Resources Create an Account Sign In

iClicker makes classroom engagement easy

Create meaningful connections with a student engagement system that works for you. Whatever your teaching style, wherever your students are, iClicker supports learning and student success with research-backed learning strategies.

Create a free instructor account

Student Preparedness Attendance Tracking Analytics & Insights Easy Integration & Setup Classroom Engagement

Choose your speed above

How can we help?

Demos & Trainings

Webinars & More

Contact Us

Key Equations

Wave speed

$$v = \frac{\lambda}{T} = \lambda f$$

Linear mass density

$$\mu = \frac{\text{mass of the string}}{\text{length of the string}}$$

Speed of a wave or pulse on a string under tension

$$|v| = \sqrt{\frac{F_T}{\mu}}$$

Speed of a compression wave in a fluid

$$v = \sqrt{\frac{B}{\rho}}$$

Resultant wave from superposition of two sinusoidal waves that are identical except for a phase shift

$$y_R(x, t) = [2A \cos\left(\frac{\phi}{2}\right)] \sin\left(kx - \omega t + \frac{\phi}{2}\right)$$

Wave number

$$k \equiv \frac{2\pi}{\lambda}$$

Wave speed

$$v = \frac{\omega}{k}$$

A periodic wave

$$y(x, t) = A \sin(kx \mp \omega t + \phi)$$

Phase of a wave

$$kx \mp \omega t + \phi$$

The linear wave equation

$$\frac{\partial^2 y(x, t)}{\partial x^2} = \frac{1}{v_w^2} \frac{\partial^2 y(x, t)}{\partial t^2}$$

Power averaged over a wavelength

$$P_{\text{ave}} = \frac{E_0}{T} \frac{1}{2} \frac{2\pi}{\lambda} \frac{1}{T} \frac{1}{2} \mu A^2 \omega^2 v$$

Intensity

$$I = \frac{P}{A}$$

Intensity for a spherical wave

$$I = \frac{P}{4\pi r^2}$$

Equation of a standing wave

$$y(x, t) = [2A \sin(kx)] \cos(\omega t)$$

Wavelength for symmetric boundary conditions

$$\lambda_n = \frac{2}{n} L, \quad n = 1, 2, 3, 4, 5, \dots$$

Frequency for symmetric boundary conditions

$$f_n = n \frac{v}{2L} = n f_1, \quad n = 1, 2, 3, 4, 5, \dots$$

A possible exam problem: (similar to online #9)

Problem requires that two formulae be put together

A guitar string of length 0.900 m is under a tension of 120.0 N. The string is made of steel. The radius of the string is 1.00 mm. What is the velocity of a transverse wave on the string?


A harder exam problem: (similar to online #9)

Problem requires that three formulae be put together

A guitar string of length 0.900 m is under a tension of 120.0 N. The string is made of steel. The radius of the string is 1.00 mm. What is the frequency of the fundamental tone?

Standing Waves

If you create a wave with fixed ends, it can only have certain wavelengths





Charge!

An electron carries the smallest possible charge.
Called “fundamental charge”

also q_e or just 'e'


$$|e| = 1.602 \times 10^{-19} \text{ Coulombs}$$

Electrons carry *negative* charge.

Protons carry the exact same *positive* charge.

Neutrons are made by crashing an electron into a proton.

Why do protons have the exact same charge as electrons?





Units1 – The SI unit of force is the:

[A] Pound

[B] gram

[C] Dyne

[D] kilogram

[E] Newton

Units2 – The SI unit of mass is the:

[A] Pound

[B] gram

[C] Dyne

[D] kilogram

[E] Newton

Units3 – The SI unit of charge is the:

[A] Volt

[B] Coulomb

[C] Ampere

[D] electron volt (eV)

[E] gram

You have 10,000 protons. How many Coulombs is this?

[A] 1 mole

[B] 10,000

[C] 1.602×10^{-19}

[D] 1.602×10^{-15}

[E] an Ampere-second

You can transfer charge to an insulator by rubbing it. (Rubbing glass with silk makes it “positive”)

You can rub a charged insulator on a Conductor to transfer charge to it.

You can charge a conductor by “induction”

You can create a force on a neutral object by polarization.



Triboseries

THESE CHARGE POSITIVE

acrylic (lucite, plexiglas)

glass

wool

silk

nylon

cotton

amber

hard rubber

saran-wrap

Teflon


THESE CHARGE NEGATIVE

(When rubbed on something higher)

Amber, or “Elektrum” from which we get “Electron” and “Electricity”



DEMOS!



DEMOS!


Soda can magic

Electroscope two types of charge

Induced Charge

Electric Forces on Metals - I

Bring a positively charged glass rod close to an electroscope without touching the sphere.




The electroscope is neutral, yet the leaves repel each other. Why?

Electric Forces on Metals - II

(b)

The electroscope is polarized by the charged rod. The sea of electrons shifts toward the rod.



Although the net charge on the electroscope is still zero, the leaves have excess positive charge and repel each other.

Modern Electricity


You can transfer charge to an insulator by spraying ions generated by a large electric field. (photocopiers / laser printers)

You can transfer charge to a conductor by connecting it to another conductor at higher voltage. (battery or power supply)


You can still do Ben Franklin things ... rubbing breaks molecular bonds

How Tribocharging Works

- *Molecular ions* can be created when one of the bonds in a large molecule is broken.
- This is the way in which a plastic rod is charged by rubbing with wool or a comb is charged by passing through your hair.




Insulator



Valence electrons are tightly bound.


Metal



Valence electrons form a "sea of electrons."


Insulators

- The electrons in an **insulator** are all tightly bound to the positive nuclei and **not free** to move around.
- Charging an insulator by friction leaves patches of molecular ions on the surface, but these patches are immobile.



Conductors

- In metals, the outer atomic electrons are only weakly bound to the nuclei.
- These outer electrons become detached from their parent nuclei and are free to wander about through the entire solid.
- The solid *as a whole* remains electrically neutral, but the electrons are now like a negatively charged liquid permeating an array of positively charged **ion cores**.



You rub a teflon with rabbits fur and make it negatively charged.

[A] The negatively charged teflon rod weighs slightly more than in did before it was rubbed and the rabbits fur a bit less.

[B] Neither material changes its weight.

[C] Both materials are lighter than before

[D] The teflon is lighter than it was

[E] Don't be silly, any weight change would be undetectable.

Which is correct?

[A] Only electrical conductors may be charged

[B] Only electrical insulators may be charged

[C] Both conductors and insulators may be charged

[D] You can't charge anything, only polarize it.

Coulomb's Law

$$F = k \frac{q_1 q_2}{r^2}$$

$$k = 8.99 \times 10^9 \frac{\text{N m}^2}{\text{C}^2}$$

What is force between electron & Proton in hydrogen?

What other law is most similar to Coulomb's Law?

- (A) Ideal gas Law
- (B) Newton's 2nd Law
- (C) Hooke's Law
- (D) Murphy's Law
- (E) Newton's Law of Gravitation

Coulomb's Law and Gravitation

$$F_E = k \frac{q_1 q_2}{r^2}$$

$$F_G = G \frac{m_1 m_2}{r^2}$$

$$k = 8.99 \times 10^9 \frac{\text{N m}^2}{\text{C}^2}$$

$$G = 6.674 \times 10^{-11} \text{N m}^2 / \text{kg}^2$$



Next Class:

More on Coulomb's law and introducing electric field