

- Announcements
  - Last assignment will be next week
  - Faraday's law and inductance (Ch 13/14)
  - Omit Ch 16
- Last Time
  - The infinite formula sheet
  - Circular motion of a charged particle
  - Motor review
  - Ampere's Law
- Today
  - Maxwell equations
  - Ampere's Law

# Drowning in Equations – Circular motion update!

An electron is accelerated through a 1000V potential in a 1 mT magnetic field. What is the radius of its circular orbit?

“Too many v’s”

$$\vec{F} = q\vec{E} + q\vec{v} \times \vec{B} \quad \text{Lorentz force Law}$$

$$U = qV \quad \text{Potential energy from potential}$$



# Maxwell's Equations

$$\oiint \vec{E} \cdot d\vec{A} = \frac{Q}{\epsilon_0} \quad \text{To calculate E for symmetrical charges.}$$

$$\oiint \vec{B} \cdot d\vec{A} = 0 \quad \text{Cannot have North magnet w/o a South pole.}$$

$$\oint \vec{B} \cdot d\vec{l} = \mu_0 I \quad \text{To calculate B for symmetrical currents.}$$

$$\oint \vec{E} \cdot d\vec{l} = -\frac{d\Phi}{dt} \quad \text{Magnetic induction! Generators! Light!}$$

$$\oiint \vec{E} \cdot d\vec{A} = \frac{Q}{\epsilon_0} \quad \text{Total flux through closed surface prop. to } Q.$$

$$\oint \vec{B} \cdot d\vec{l} = \mu_0 I \quad \text{Line integral of } B \text{ prop. to current thru loop.}$$

## **Electric fields are calculated with Coulomb's law or Gauss's Law**

$$\vec{E} = \frac{1}{4\pi\epsilon_0} \frac{Q\hat{r}}{r^2} \quad \text{Coulomb}$$

$$\oiint \vec{E} \cdot d\vec{A} = \frac{Q_{\text{enclosed}}}{\epsilon_0} \quad \text{Gauss's Law}$$

## **Magnetic Fields are calculated with the Biot-Savart Law or Ampere's Law**

$$d\vec{B} = \frac{\mu_0}{4\pi} \frac{I d\vec{l} \times \hat{r}}{r^2} \quad \text{Biot Savart}$$



$$\oint \vec{B} \cdot d\vec{l} = \mu_0 I_{\text{enclosed}} \quad \text{Ampere's Law.}$$

$$\oiint \vec{E} \cdot d\vec{A} = \frac{Q_{\text{enclosed}}}{\epsilon_0}$$

Gauss to calculate E for symmetrical charges.

$$\vec{E} = \frac{\sigma}{2\epsilon_0} \hat{k}$$



Planar symmetry



Infinite parallel-plate capacitor

$$\vec{E} = \frac{\lambda}{2\pi r \epsilon_0} \hat{r}$$



Cylindrical symmetry



Coaxial cylinders

$$\vec{E} = \frac{Q}{4\pi r^2 \epsilon_0} \hat{r}$$

Spherical symmetry



Concentric spheres

# Equations of Magnetism

$$\vec{F} = Q \vec{v} \times \vec{B} \quad \text{Force on charge } Q$$

$$\vec{F} = I \vec{L} \times \vec{B} \quad \text{Force on current } I$$

$$\vec{B} = \frac{\mu_0 I}{2\pi r} \hat{\phi} \quad \text{Field of Infinite wire}$$

$$\vec{B} = \frac{\mu_0 I r}{2\pi R^2} \hat{\phi} \quad \text{Field inside a wire of radius } R$$

$$\vec{B} = \mu_0 n I \hat{z} \quad \text{Field of an infinite coil (solenoid)}$$

$$\vec{B} = \frac{\mu_0 I}{2a} \hat{z} \quad \text{Field in center of wire loop}$$





# Equations from Ampere's Law

$$\vec{B} = \frac{\mu_0 I}{2\pi r} \hat{\phi} \quad \text{Field of Infinite wire}$$

$$\vec{B} = \frac{\mu_0 I r}{2\pi R^2} \hat{\phi} \quad \text{Field inside a wire of radius } R$$

$$\vec{B} = \mu_0 n I \hat{z} \quad \text{Field of an infinite coil (solenoid)}$$

# Advanced Math Alert!! (Optional)




$$\oiint \vec{E} \cdot d\vec{A} = \frac{Q_{\text{enclosed}}}{\epsilon_0}$$


What's a surface integral really?

$\oint \vec{F}(u, v) \cdot \hat{n} \, dA$  Vector closed surface integral


$$\oint \vec{E} \cdot d\vec{A} = \frac{Q}{\epsilon_0}$$



$$\int f(x, y) dA = \iint f(x, y) dx dy \quad \text{Volume under an area}$$




$$\int f(u, v) dA = \iint f(u, v) |\vec{r}_u \times \vec{r}_v| du dv \quad \text{Surface integral}$$




$\int \vec{F}(u, v) \cdot \hat{n} \, dA$  Vector surface integral

$$\iint \vec{F} \cdot (\vec{r}_u \times \vec{r}_v) \, du \, dv$$



$\oint \vec{F}(u, v) \cdot \hat{n} \, dA$  Vector closed surface integral

$$\oint \vec{E} \cdot d\vec{A} = \frac{Q}{\epsilon_0}$$






$$\oiint \vec{E} \cdot d\vec{A} = \frac{Q_{\text{enclosed}}}{\epsilon_0}$$

Gauss to calculate E for symmetrical charges.


$$\vec{E} = \frac{\sigma}{2\epsilon_0} \hat{k}$$

Planar symmetry




$$\vec{E} = \frac{\lambda}{2\pi r \epsilon_0} \hat{r}$$

Cylindrical symmetry




$$\vec{E} = \frac{Q}{4\pi r^2 \epsilon_0} \hat{r}$$


Spherical symmetry



Infinite parallel-plate capacitor



Coaxial cylinders



Concentric spheres


# Equations from Ampere's Law

$$\vec{B} = \frac{\mu_0 I}{2\pi r} \hat{\phi} \quad \text{Field of Infinite wire}$$

$$\vec{B} = \frac{\mu_0 I r}{2\pi R^2} \hat{\phi} \quad \text{Field inside a wire of radius R}$$

$$\vec{B} = \mu_0 n I \hat{z} \quad \text{Field of an infinite coil (solenoid)}$$

$$\oint \vec{B} \cdot d\vec{l} = \mu_0 I_{\text{enclosed}} \quad \text{Ampere's Law.}$$



Given  $i_1 = 10$  Amps,  $i_2 = -5$  A,  $i_3 = 5$  A


What is  $\int \vec{B} \cdot d\vec{l}$ ?

(A) Plan 9 from outer space

(B)  $10\mu_0 \text{ T}\cdot\text{m}$

(C)  $15\mu_0 \text{ T}\cdot\text{m}$


(D)  $5\mu_0 \text{ T}\cdot\text{m}$






# Ampere applied inside and outside of a wire

All of the current is encircled and thus all is used in Ampere's law.





Only the current encircled by the loop is used in Ampere's law.



# Ampere applied outside of a wire

All of the current is encircled and thus all is used in Ampere's law.





$$\vec{B} = \frac{\mu_0 I}{2\pi r} \hat{\phi}$$

Field outside long wire





# Ampere applied inside of a wire




$$\vec{B} = \frac{\mu_0 I r}{2 \pi a^2} \hat{\phi}$$

Field inside a wire of radius  $a$

**A wire with radius 2 cm carries 8 Amperes  
How many Amps are contained within 1 cm  
radius?**

- (A)** 16
- (B)** 8
- (C)** 4
- (D)** 2




**A wire with radius “a” carries “I” Amperes  
How many Amps are contained within  
radius “r”?**

**(A)**  $I$

**(B)**  $\frac{I}{4}$

**(C)**  $I \frac{r}{a}$

**(D)**  $I \frac{r^2}{a^2}$





# Sanity Check: What happens at $r = R$ ?




# What happens at $r = R$ ?


$$\vec{B} = \frac{\mu_0 I r}{2 \pi R^2} \hat{\phi} \quad \text{Field inside a wire of radius } R$$

$$\vec{B} = \frac{\mu_0 I}{2 \pi r} \hat{\phi} \quad \text{Field outside long wire}$$

# Ampere result inside and outside of a wire




# Derive field of a solenoid



$\vec{B} = \mu_0 n I \hat{z}$  Field of an infinite coil (solenoid)









$\vec{B} = \mu_0 n I \hat{z}$     Field of an infinite coil (solenoid)    33

If the solenoid carries current “I” and there are “N” windings inside the amperian loop what is  $\oint \vec{B} \cdot d\vec{l}$ ?



- (A) I
- (B)  $\mu_0 N I$
- (C)  $\mu_0 I/N$
- (D)  $-\mu_0 I$



$\vec{B} = \mu_0 n I \hat{z}$     Field of an infinite coil (solenoid)    35

$$\vec{B} = \mu_0 n I \hat{z} \quad \text{Field of an infinite coil (solenoid)}_{36}$$

If the solenoid carries current “I” and there are “N” windings inside the amperian loop what is  $\oint \vec{B} \cdot d\vec{l}$ ?

- (A) I
- (B)  $\mu_0 N I$
- (C)  $\mu_0 I/N$
- (D)  $-\mu_0 I$

**If the coil shown is 10 cm long and has a total of 4000 windings and carries two Amperes, what is the magnetic field inside?**



- (A)** 0.25 T
- (B)** 0.025 T
- (C)** 1 T
- (D)** 0.1 T

$$\vec{B} = \mu_0 n I \hat{z}$$







# Ferromagnets, Permeability and Susceptibility

$\vec{B} = \mu_0 n I \hat{z}$     Field of an infinite coil (solenoid)

$\vec{B} = \mu_0 (1 + \chi) n I \hat{z}$     With susceptibility

$\vec{B} = \mu n I \hat{z}$     With permeability  $\mu$



# Recap Lecture 24

- Amperes law
- Field of a wire
- Field of a solenoid
-

# Maxwell's Equations

$$\oiint \vec{E} \cdot d\vec{A} = \frac{Q_{\text{enclosed}}}{\epsilon_0} \quad \text{To calculate E for symmetrical charges.}$$

$$\oiint \vec{B} \cdot d\vec{A} = 0 \quad \text{Cannot have North magnet w/o a South pole.}$$

$$\oint \vec{B} \cdot d\vec{l} = \mu_0 I_{\text{enclosed}} \quad \text{To calculate B for symmetrical currents.}$$

$$\oint \vec{E} \cdot d\vec{l} = -\frac{d\Phi}{dt} \quad \text{Magnetic induction! Generators! Light!}$$

$$\Phi_B = \int \vec{B} \cdot d\vec{A}$$

# How do I avoid an infinite formula sheet – really?

$$\Delta V = - \int \vec{E} \cdot d\vec{l} \quad \text{Definition of potential.}$$

$$\oint \vec{E} \cdot d\vec{A} = \frac{Q}{\epsilon_0} \quad \text{Gauss Law (sphere/line/plane).}$$

$$Q = CV \quad \text{Def. of capacitance}$$

$$V = IR \quad R = \rho \frac{L}{A} \quad \text{Ohm's Law and origin of resistance}$$

$$U = qV \quad \text{Relation between potential and potential energy}$$

$$\vec{F} = q\vec{E} + q\vec{v} \times \vec{B} \quad \text{Lorentz force Law}$$

$$\oint \vec{B} \cdot d\vec{l} = \mu_0 I \quad \text{Ampere's law (wire/solenoid/current sheet).}$$